arXiv:1108.3215v1 [hep-th] 16 Aug 2011

Gauge Invariant Cutoff QED

F. Ardalan[a,b],[*] H. Arfaei[a,c],[†] M. Ghasemkhani[c],[‡] and N. Sadooghi[a§]
[a]*Department of Physics, Sharif University of Technology, P.O. Box 11155-9161, Tehran-Iran*
[b]*Institute for Research in Fundamental Sciences (IPM)*
School of Physics, P.O. Box 19395-5531, Tehran-Iran
[c]*Institute for Research in Fundamental Sciences (IPM)*
School of Particles and Accelerators, P.O. Box 19395-5531, Tehran-Iran

Abstract

A hidden generalized gauge symmetry of a cutoff QED is used to show the renormalizability of QED. In particular, it is shown that corresponding Ward identities are valid all along the renormalization group flow. We also derive the corresponding exact RG flow equation of the effective average action for a cutoff $\lambda\varphi^4$ theory. Generalization to any gauge group is indicated.

PACS numbers: 11.10.Gh, 11.30.-j, 11.10.Hi

I. INTRODUCTION

Diagrammatic proof of the renormalizability of QED in the BPHZ context is a lengthy derivation [1]. In 1984, Polchinski presented a simple proof using a version of Wilsonian renormalization group equation (RGE) for the $\lambda\varphi^4$ theory [2]. Although Polchinski's proof of renormalizability could in principle be extended to QED, the manifest violation of gauge invariance in his momentum cutoff formulation was an obstacle to a straightforward proof. This has been extensively studied and is by now a well understood derivation of renormalizability of QED [3, 4] (for recent reviews see [5] and the literature therein). Yet, it remains an untidy procedure involving either complicated cutoff insertions or additional labor for the proof of Ward identities.

In this work, we present a simple extension of Polchinski's proof of renormalizability to QED using a hidden generalized gauge invariance of the cutoff formulation, thus circumventing the need for explicit verification of the Ward identities. In Sec. II, we introduce our slightly modified cutoff procedure of Polchinski and apply it to the $\lambda\varphi^4$ theory. The cutoff procedure we use is simply multiplying the fields by a cutoff function in the momentum representation, as explained in Sec. II. This procedure has its origin in Kogut and Wilson's [6] "Incomplete Integration", with which Wilson's exact RGE was first formulated later. Wetterich introduced a similar concept of "average field" in the discussion of RGE [7], which was a precursor to his well-known presentation of exact average RGE [5, 8–10]. We arrived

---

[*] Electronic address: ardalan@ipm.ir
[†] Electronic address: arfaei@ipm.ir
[‡] Electronic address: ghasemkhani@ipm.ir
[§] Electronic address: sadooghi@physics.sharif.ir

at this cutoff procedure from an entirely different angle of translationally invariant noncommutative gauge theory [13, 14], which led us to the symmetry of the cutoff effective action for gauge theories, explained in Sec. III. Here, we present the generalized gauge symmetry of the cutoff QED, and indicate its renormalizability. The cutoff procedure we use and the consequent symmetry can be easily generalized to the gauge theory with any group and matter field, including $SU(N)$, as the noncommutative structure from which it is derived in Sec. III can be extended trivially to these cases. The point being that the "noncommutative geometric" structure we will be using, is in fact Abelian. For simplicity of presentation, we will restrict ourselves to the case of $U(1)$ gauge group, QED. In Sec. IV, we derive the corresponding exact RG flow equation for the effective average action of $\lambda\varphi^4$ theory using an IR cutoff function, a method developed in [8, 9]. We will show that it is different from the one derived there. The main reason for this difference is the multiplicative nature of our cutoff function, in contrast to the additive IR cutoff introduced in [8, 9]. Modifying each field with such a multiplicative cutoff, the cutoff function appears not only in the kinetic term, but also in the interacting part of the classical cut off action. It is interesting to generalize this procedure to gauge theories and to explore the possible practical consequences of the new exact average RG for the effective average action as well as the hidden generalized gauge symmetry along the flow, pointed out in this paper. The origin of this gauge symmetry will nevertheless be discussed in the conclusions, Sec. V, and Appendix A.

II. MODIFIED CUTOFF REGULARIZATION

In this section we redo Polchinski's proof of $\lambda\varphi^4$ theory [2] by a slightly different cutoff procedure. The real scalar theory defined by the action

$$S = \int d^4x\; [-\frac{1}{2}\partial_\mu\varphi\partial^\mu\varphi + \frac{1}{2}m^2\varphi^2 + \frac{1}{4!}\lambda\varphi^4], \tag{II.1}$$

and the momentum cutoff Λ_0 in the Euclidean space is regularized by Polchinski via introduction of a momentum cutoff $K_{\Lambda_0}(p)$ into the propagator,

$$\frac{K_{\Lambda_0}(p)}{p^2+m^2}. \tag{II.2}$$

The main property of the function $K_{\Lambda_0}(p)$ is that it has a value equal to 1 for $p^2 < \Lambda_0^2$ and decreases rapidly for $\frac{p^2}{\Lambda_0^2} \to \infty$. An example is

$$K_{\Lambda_0}(p) \;=\; \begin{cases} 1, & p^2 \leq \frac{\Lambda_0^2}{4}, \\ \exp[(1-\frac{p^2}{\Lambda_0^2})^{-1}\exp(4-\frac{p^2}{\Lambda_0^2})^{-1}], & \frac{\Lambda_0^2}{4} < p^2 < \Lambda_0^2, \\ 0, & p^2 \geq \Lambda_0^2. \end{cases} \tag{II.3}$$

The cutoff function $K_{\Lambda_0}(p)$ effectively cuts off the momentum integral in all loops, rendering them ultraviolet finite in perturbation theory. The introduction of a cutoff function in the propagators, and consequently in all loops is an efficient procedure of implementing Wilson's renormalization group flow, as the cutoff momentum p can now be lowered all the way down to zero in the path integral for the effective action. The remarkable aspect of the introduction of the cutoff function in the loop integrals is that loop integrals can now be estimated easily and cutoff independence of Green's functions be demonstrated in a few steps in marked contrast to the lengthy BPHZ proof of renormalizability.

In this work, we propose to modify this procedure by extending the cutoff to all the terms in the action, and in fact to all fields. It must be immediately emphasized that multiplication of fields by a cutoff function is not a trivial field redefinition when the cutoff function vanishes for momenta larger than Λ. In our formulation we replace the field $\tilde{\varphi}(p)$ by their cutoff counterpart

$$h_\Lambda(p)\tilde{\varphi}(p), \tag{II.4}$$

in momentum space. Here, the cutoff functions are assumed to vanish for $p^2 > \Lambda^2$

$$h_\Lambda(p) = 0, \qquad p^2 > \Lambda^2. \tag{II.5}$$

This is therefore a straightforward implementation of Wilson's cutoff procedure executed directly on the fields rather than on the path integrals.

In the original Polchinski's formulation, cutoff independence of the theory is

$$\frac{\partial \mathcal{Z}}{\partial t} = 0, \qquad \text{with} \qquad \frac{\partial}{\partial t} \equiv \Lambda \frac{\partial}{\partial \Lambda}, \tag{II.6}$$

and, the running effective action is

$$S_{\text{eff}} = \int \frac{d^4p}{(2\pi)^4}\Big[-\frac{1}{2}\tilde{\varphi}(p)\frac{p^2+m^2}{K_\Lambda(p)}\tilde{\varphi}(-p) + \mathcal{L}_{\text{int}}(\tilde{\varphi},\Lambda)\Big]. \tag{II.7}$$

Equation (II.6) then determines the running effective interaction lagrangian $\mathcal{L}_{\text{int}}$ at the Λ scale, which now satisfies the functional differential equation

$$\frac{\partial \mathcal{L}_{\text{int}}}{\partial t} = -\frac{1}{2}\int \frac{d^4p}{(2\pi)^4}\frac{1}{p^2+m^2}\,\frac{\partial K_\Lambda(p)}{\partial t}\Big[\frac{\partial \mathcal{L}_{\text{int}}}{\partial \tilde{\varphi}(p)}\frac{\partial \mathcal{L}_{\text{int}}}{\partial \tilde{\varphi}(-p)} + \frac{\partial^2 \mathcal{L}_{\text{int}}}{\partial \tilde{\varphi}(p)\partial \tilde{\varphi}(-p)}\Big]. \tag{II.8}$$

Our formulation does not yet differ from Polchinski's as the functional $\mathcal{L}_{\text{int}}$ is not yet specified. The difference shows up when expanding $\mathcal{L}_{\text{int}}$ in a series in $\tilde{\varphi}$.

$$\mathcal{L}_{\text{int}}(\tilde{\varphi},\Lambda) = \sum_{n=1}^{\infty}\frac{1}{(2n)!}\int \frac{d^4p_1\cdots d^4p_{2n}}{(2\pi)^{8n-4}}\; L_{2n}(p_1,\cdots,p_{2n};\Lambda)\delta(\sum_i p_i)\;\tilde{\varphi}(p_1)\cdots\tilde{\varphi}(p_{2n}). \tag{II.9}$$

In Polchinski's formulation the renormalization flow equation (II.8) reduces to a set of equations for the coefficient functions $L_{2n}(p_1,\cdots,p_{2n},\Lambda)$, whose solution would give the effective action at scale Λ,

$$\begin{aligned}
&(\frac{\partial}{\partial t} + 4 - 2n)L_{2n}(p_1,\cdots,p_{2n};\Lambda)\\
&= -\sum_{\ell=1}^{n}\Big\{Q_\Lambda(P,m)L_{2\ell}(p_1,\cdots,p_{2\ell-1};\Lambda)L_{2n+2-2\ell}(p_{2\ell},\cdots,p_{2n},-P;\Lambda) + permutation\\
&\qquad -\frac{1}{2}\int\frac{d^4p}{(2\pi\Lambda)^4}L_{2n+2}(p_1,\cdots,p_{2n},P,-P;\Lambda)Q_\Lambda(P,m)\Big\}.
\end{aligned} \tag{II.10}$$

Here, $P = \sum_{i=1}^{2\ell-1} p_i$, and

$$Q_\Lambda(P,m) = \frac{1}{P^2+m^2}\,\Lambda^2\frac{\partial K_\Lambda(P)}{\partial t}. \tag{II.11}$$

A convenient rescaling $L_{2n} \to \Lambda^{4-2n}L_{2n}$ has been inserted in (II.10).

In our formulation, however, the expansion in (II.9) has to be replaced by

$$\mathcal{L}_{\text{int}}(\tilde{\varphi},\Lambda)=\sum_{n=1}^{\infty}\frac{1}{(2n)!}\int\frac{d^4p_1\cdots d^4p_{2n}}{(2\pi)^{8n-4}}\;L'_{2n}(p_1,\cdots,p_{2n};\Lambda)\delta(\sum_i p_i)\;h_\Lambda(p_1)\tilde{\varphi}(p_1)\cdots h_\Lambda(p_{2n})\tilde{\varphi}(p_{2n}),\tag{II.12}$$

as every momentum space field $\tilde{\varphi}(p)$ is rescaled and cut off by the cutoff function $h_\Lambda(p)$ from (II.4). It may seem that the subsequent steps in renormalizality may get complicated, but this is not the case. In fact the resulting renormalization group flow equations for the function $h_\Lambda(p_1)\cdots h_\Lambda(p_{2n})L'_{2n}(p_1,\cdots,p_{2n};\Lambda)$, now would be exactly the same form as (II.10) for the coefficient functions, where the function $K_\Lambda(p)$ of Polchinski has to be replaced by $h_\Lambda^{-2}(p)$, see [2] for more details. The advantage of Polchinski's approach to renormalization is that the insertion of a cutoff function in the integrals allows him to estimate the coefficient functions $L_{2n}(p_1,\cdots,p_{2n})$ through functional analytic methods and put bounds on them; and then to prove cutoff independence of the Green's functions in perturbation theory, through a series of Lemmas. Note that in all this the function K_Λ appears, as in (II.8), only in its derivative form $\partial_t K_\Lambda$, which does not affect the estimate arguments which involve various norms of functions. We will not go through the entire analysis, but only point out the validity of the procedure in our approach in cases where there is ground for doubt.

The first instance that our approach may invalidate Polchinski's result is when a bound on $L'_{2n}(p_1,\cdots,p_{2n})$ is obtained through (II.10) using the bounds

$$\int\frac{d^4p}{(2\pi)^4}|Q_\Lambda(p,m)|<C\Lambda^4,\tag{II.13}$$

and

$$\max\left|\frac{\partial^n}{\partial p^n}Q_\Lambda(p,m)\right|<D_n\Lambda^{-n}.\tag{II.14}$$

Here, C and D_n are appropriate constants. The result

$$\begin{aligned}\max\left|L_{2n}^{(r)}(p_1,...,p_{2n};\Lambda)\right| &\leq \mathcal{P}_{2r-n}(\ln\tfrac{\Lambda_0}{\Lambda}), &&\text{for}\quad r+1-n\geq 0,\\ &= 0, &&\text{for}\quad r+1-n<0,\end{aligned}\tag{II.15}$$

and similar bounds are central to the proof of renormalizability. In (II.15), $L_{2n}^{(r)}$ is the r'th term in the perturbative expansion of L_{2n}, and $\mathcal{P}_{2r-n}$ are polynomials of order $2r-n$.

In our case, (II.10) involves $h_\Lambda(p_1)\cdots h_\Lambda(p_n)L_{2n}$ rather than L_{2n}. One must make sure that the appearance of the h_Λ's does not ruin the bounds in Polchinski and therefore ruin the arguments on renormalizability of the theory. We have identified the function K_Λ of (II.2) with h_Λ^{-2} of (II.4). Therefore if K_Λ's should behave as in Polchinski's formulation, i.e. go to zero when p^2 approaches Λ^2, and vanish for $p^2>\Lambda^2$, then h_Λ should become large as p^2 approaches Λ^2. Of course, we set $h_\Lambda(p^2)=0$ for $p^2>\Lambda^2$. Therefore in the left hand side (l.h.s.) of (II.10), after inserting $h_\Lambda\cdots h_\Lambda L'_{2n}$ for L_{2n}'s, as the h_Λ's are larger than one in their range of definition, they can be dropped in the ensuing inequality. On the right hand side (r.h.s.) of (II.10), we now use the functional norm with an appropriate weight to kill off the value of $h_\Lambda>1$ for $p^2\to\Lambda^2$,[1] leading to the desired inequality (II.15), this time for L'_{2n}'s.

---

[1] See footnote on page 280 of Polchinski [2].

III. GAUGE INVARIANCE OF CUTOFF QED AND WARD IDENTITIES

The Polchinski's procedure was applied to QED very early and provided a simple proof of renormalizability of the theory [3]. In QED there are now two propagators to cut off, that of the electron and that of the photon. This can be done with the same cutoff function. The renormalization group equation is similarly derived and estimates for the coefficient functions and bounds on Green's functions obtained. The situation is then a straightforward extension of the $\lambda\varphi^4$ theory. There is only one significant hurdle to overcome which has engaged authors of Refs. [3, 4] ever since and is the subject of the present work, *i.e.* the question of gauge invariance.

The problem is that Polchinski's approach, and in fact any approach involving a momentum cutoff, inherently violates gauge invariance: Gauge invariance is a statement about the behavior of gauge fields in a space-time point, involving all momenta. Thus, at any finite cutoff scale Λ, the flow equation and its solutions are not gauge invariant. However, it was proved, that the final IR point of the flow $\Lambda \to 0$, the expressions for the quantum effective action and the Green's functions are indeed invariant [4]. There were also nontrivial modifications of the cutoff procedure which were not gauge invariant all along the flow, but only at its end points. These formulations have been extensively pursued in the application of the exact renormalization group in such areas as QCD (see [5] for recent reviews), and gravity [12]. In all these works derivation of the modified Ward-Takahashi identities is the essential complication.

In this section, we will show that our version of introduction of the momentum cutoff in the theory ensures persistence of gauge invariance in the form of a generalized symmetry of the cutoff QED and derive the resultant Ward-Takahashi identities in a standard manner. We start from the classical action of QED

$$S_{\text{QED}} = \int d^4x \Big\{ -\bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi + e\bar{\psi}\gamma^\mu A_\mu \psi + \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2\xi}(\partial_\mu A^\mu)^2 \Big\}, \tag{III.1}$$

with the cutoff function $h_\Lambda(p)$ inserted on each field in the momentum space

$$\tilde{\psi}(p) \to h_\Lambda(p)\tilde{\psi}(p), \qquad \text{and} \qquad \tilde{A}_\mu(p) \to h_\Lambda(p)\tilde{A}_\mu(p). \tag{III.2}$$

Here, the cutoff function $h_\Lambda(p)$ has the property of being equal to 1 for $p^2 < \Lambda^2$ and $h_\Lambda(p) \to 0$ rapidly for $p^2 > \Lambda^2$. We denote this cutoff action as S_{h_Λ}. This action has a symmetry which is the generalization of gauge symmetry of S_{QED}. Whereas the gauge symmetry of S_{QED} is

$$\psi(x) \to e^{ie\epsilon(x)}\psi(x), \qquad \bar{\psi}(x) \to e^{-ie\epsilon(x)}\bar{\psi}(x), \qquad A_\mu \to A_\mu(x) - \partial_\mu\epsilon(x), \tag{III.3}$$

the symmetry of S_{h_Λ} is similarly defined but now involves h_Λ. First note that when in momentum space two functions $\tilde{\psi}(p)$ and $h_\Lambda(p)$ are pointwise multiplied, their corresponding functions in the configuration space, $\psi(x)$ and $\tilde{h}_\Lambda(x)$, are multiplied via convolution,

$$h_\Lambda(p)\tilde{\psi}(p) \to \tilde{h}_\Lambda(x) \circ \psi(x), \tag{III.4}$$

where convolution of two functions $f(x)$ and $g(x)$ is defined by,

$$(f \circ g)(x) \equiv \int d^4y \; f(x-y)g(y). \tag{III.5}$$

The above mentioned gauge symmetry transformation of S_{h_Λ} is

$$\psi(x) \to \widetilde{(h_\Lambda^{-1})} \circ [(\tilde{h}_\Lambda \circ g)(\tilde{h}_\Lambda \circ \psi)]. \tag{III.6}$$

Here, $g(x)$ is the generalization of $e^{ie\epsilon(x)}$ defined by

$$g(x) = 1 + ie\epsilon(x) + \frac{1}{2!}\widetilde{(h_\Lambda^{-1})} \circ [(\tilde{h}_\Lambda \circ (ie\epsilon))(\tilde{h}_\Lambda \circ (ie\epsilon))] + \cdots \tag{III.7}$$

The transformation of $A_\mu(x)$ is

$$A_\mu(x) \to A_\mu(x) + \widetilde{(h_\Lambda^{-1})} \circ \Big[\Big(\tilde{h}_\Lambda \circ g\Big) \Big(\tilde{h}_\Lambda \circ (\partial_\mu g^{-1})\Big) \Big]. \tag{III.8}$$

These strange looking transformations come from a simple generalized noncommutative geometric construction described in the Appendix A.

We note that the relations (III.6) and (III.8), which are not local anymore, reduce to the usual gauge transformations of QED when $h_\Lambda = 1$. These generalized "cutoff gauge transformations" are shown in the Appendix A to be the symmetry of S_{h_Λ}. They ensure the "good" properties of the effective action S_{h_Λ} all along the renormalization group flow, including transversality of the photon, and exclusion of unwanted non-gauge invariant terms in the effective action S_{h_Λ}.

We will now write down the consequent Ward-Takahashi identities, with the derivations described in Appendix B. There are essentially two different well-known such identities which take the following forms in our case,

$$\begin{aligned}
&-\frac{1}{\xi}p^2 p_\mu \frac{\delta W}{\delta \tilde{J}_\mu(-p)} + h_\Lambda^2(p) p_\mu \tilde{J}^\mu(p) \\
&\qquad -eh_\Lambda(p)\int \frac{d^4q}{(2\pi)^4}\left[h_\Lambda(q)h_\Lambda^{-1}(q-p)\frac{\delta W}{\delta\tilde{\chi}(q-p)}\tilde{\chi}(q) + h_\Lambda^{-1}(q)h_\Lambda(q-p)\tilde{\bar{\chi}}(q-p)\frac{\delta W}{\delta\tilde{\bar{\chi}}(q)}\right] = 0,
\end{aligned} \tag{III.9}$$

for the generating function $W[J_\mu, \chi, \bar{\chi}]$ of connected Green's function, and

$$\begin{aligned}
&-\frac{1}{\xi}p^2 p_\mu h_\Lambda^2(p) A^\mu(p) - p_\mu \frac{\delta\Gamma}{\delta\tilde{A}_\mu(-p)} \\
&\qquad +eh_\Lambda(p)\int \frac{d^4q}{(2\pi)^4}\left[h_\Lambda^{-1}(q)h_\Lambda(q-p)\tilde{\bar{\psi}}(q-p)\frac{\delta\Gamma}{\delta\tilde{\bar{\psi}}(q)} + h_\Lambda(q)h_\Lambda^{-1}(q-p)\frac{\delta\Gamma}{\delta\tilde{\psi}(q-p)}\tilde{\psi}(q)\right] = 0,
\end{aligned} \tag{III.10}$$

for the generating functional $\Gamma[A_\mu, \bar{\psi}, \psi]$ one particle irreducible (1PI) vertex functions. It is known that $W[J_\mu, \chi, \bar{\chi}]$ and $\Gamma[A_\mu, \bar{\psi}, \psi]$ are related through the Legendre transformation

$$\Gamma[\tilde{A}_\mu, \tilde{\bar{\psi}}, \tilde{\psi}] = W[\tilde{J}_\mu, \tilde{\chi}, \tilde{\bar{\chi}}] - \int \frac{d^4p}{(2\pi)^4} h_\Lambda^2(p)\Big[\tilde{\bar{\chi}}(p)\bar{\psi}(p) + \tilde{\bar{\psi}}(p)\tilde{\chi}(p) + \tilde{J}_\mu(p)\tilde{A}^\mu(-p)\Big], \tag{III.11}$$

with respect to the sources $(\tilde{J}_\mu, \tilde{\chi}, \tilde{\bar{\chi}})$ corresponding to the fields $(\tilde{A}_\mu, \tilde{\bar{\psi}}, \tilde{\psi})$ in momentum space. The Ward identities (III.9) and (III.10) correspond to the usual Ward identities of QED with identical physical significance establishing the rôle of our generalized "cutoff gauge invariance" in the renormalization group flow and the renormalization program of QED.

IV. EXACT RG FLOW EQUATION FOR THE EFFECTIVE AVERAGE ACTION OF CUTOFF $\lambda\varphi^4$ THEORY

In a separate development an alternative renormalization group equation was derived for the effective average action [5, 8–10] to be defined below. The idea was to add an IR cutoff term

$$\Delta\mathcal{S}_k[\varphi] = \frac{1}{2}\int \frac{d^4q}{(2\pi)^4}\mathcal{R}_k(q)\tilde{\varphi}(-q)\tilde{\varphi}(q), \tag{IV.1}$$

to the classical action (II.1) in the Euclidean space, and to modify in this way the standard effective action of the theory. In (IV.1), the IR cutoff $\mathcal{R}_k$ satisfies the following properties

$$\mathcal{R}_k(q)\begin{cases} = 0 & \text{for } k \to 0 \\ \to \infty & \text{for } k \to \Lambda \text{ or } k \to \infty. \end{cases} \tag{IV.2}$$

An example for $\mathcal{R}_k(q)$, which is also used in [10] is

$$\mathcal{R}_k(q) \sim \frac{q^2}{e^{\frac{q^2}{k^2}} - 1}, \tag{IV.3}$$

that behaves as $\mathcal{R}_k(q) \sim k^2$ for fluctuations with small momenta $q^2 \ll k^2$, and vanishes for $q^2 \gg k^2$. Adding $\Delta\mathcal{S}_k[\varphi]$ to the classical action and integrating over all fluctuations to derive the effective action of the theory will induce automatically an effective mass $\sim k$ to those Fourier modes of $\tilde{\varphi}(q)$ with small momenta $q^2 \ll k^2$, and prohibit the high momentum modes with $q^2 \gg k^2$ from contributing in the effective action of the theory. The resulting effective average action $\Gamma_k[\phi]$ will depend on the scale k and satisfies the RG flow equation [see e.g. [5, 8–10] for a rigorous derivation of (IV.4)]

$$\partial_t\Gamma_k[\phi] = \frac{1}{2}\text{Tr}\Big\{G_k^{(2)}\partial_t\mathcal{R}_k\Big\}, \tag{IV.4}$$

where $\phi \equiv \langle\varphi\rangle$ and $\partial_t \equiv k\partial_k$. The trace involves an integration over momenta. Moreover, $G_k^{(2)}$ is the full connected two-point Green's function satisfying

$$G_k^{(2)} = [\Gamma_k^{(2)} + \mathcal{R}_k]^{-1}. \tag{IV.5}$$

Here, $\Gamma_k^{(2)}$ is the exact of 1PI two-point vertex function, arising from variation of the effective average action $\Gamma_k[\phi]$ two times with respect to ϕ. By definition, the effective average action interpolates between the classical action, $\Gamma_\Lambda \approx S_0$, and the full effective action $\Gamma = \lim\limits_{k\to 0}\Gamma_k$ [10]. In Γ_Λ, Λ is a natural cutoff that characterizes the theory. The diagrammatic representation of (IV.4) is presented in Fig. 1.

In this section, we will present the exact flow equation of the effective average action of a cutoff $\lambda\varphi^4$ theory with our cutoff procedure of (II.4). Our goal is to compare the final form of the corresponding RG flow equation with (IV.4). Applications of the new RG flow equation and the consequences of the new hidden gauge invariance, pointed out in Sec. III, will be presented elsewhere [11]. To derive the above mentioned flow equation, let us start by considering the action (II.1) in the Euclidean space and to replace all fields $\tilde{\varphi}(q)$ in momentum space by $h_k(q)\tilde{\varphi}(q)$. In contrast to h_Λ in (II.4), $h_k(q)$ is



$$\partial_t\Gamma_k[\phi] = \tfrac{1}{2}$$

FIG. 1: Diagramatic representation of the exact RG flow equation (IV.4). The thick black line represents the full connected two-point Green's function in the presence of the additional IR cutoff $\mathcal{R}_k(q)$, i.e. $G_k^{(2)} = [\Gamma_k^{(2)} + \mathcal{R}_k]^{-1}$. The filled green box represents the insertion of a factor $\partial_t\mathcal{R}_k$.

assumed to depend only on the momentum q and the RG scale k and not on the cutoff Λ. With this replacement, the modified cutoff action then reads

$$\begin{aligned} S_k[\varphi] &= \frac{1}{2}\int \frac{d^4q_1}{(2\pi)^4}\frac{d^4q_2}{(2\pi)^4}\mathcal{H}_k^{(2)}(q_1,q_2)\tilde{\varphi}(q_1)\tilde{\varphi}(q_2) \\ &\qquad + \int \frac{d^4q_1}{(2\pi)^4}\cdots\frac{d^4q_4}{(2\pi)^4}\mathcal{H}_k^{(4)}(q_1,\cdots,q_4)\tilde{\varphi}(q_1)\tilde{\varphi}(q_2)\tilde{\varphi}(q_3)\tilde{\varphi}(q_4), \end{aligned} \tag{IV.6}$$

where the cutoff functions $\mathcal{H}_k^{(2)}$ and $\mathcal{H}_k^{(4)}$ are defined by

$$\begin{aligned} \mathcal{H}_k^{(2)}(q_1,q_2) &\equiv h_k(q_1)h_k(q_2)(q_1^2+m^2)\delta(q_1+q_2), \\ \mathcal{H}_k^{(4)}(q_1,q_2,q_3,q_4) &\equiv \frac{\lambda}{4!}h_k(q_1)h_k(q_2)h_k(q_3)h_k(q_4)\delta(q_1+q_2+q_3+q_4). \end{aligned} \tag{IV.7}$$

In App. C, we will follow the method described in [8],[2] and will derive the corresponding exact RG flow equation to the action (IV.6). We will show that the RG flow equation of the effective average action of the cutoff $\lambda\varphi^4$ theory is given by [see also (C.11)]

$$\begin{aligned} \frac{\partial\Gamma_k[\phi]}{\partial t} &= \frac{1}{2}\int \frac{d^4q_1}{(2\pi)^4}\frac{d^4q_2}{(2\pi)^4}\frac{\partial\mathcal{H}_k^{(2)}(q_1,q_2)}{\partial t}[G_k^{(2)}(q_1,q_2)+\tilde{\phi}(q_1)\tilde{\phi}(q_2)] \\ &+ \int \frac{d^4q_1}{(2\pi)^4}\cdots\frac{d^4q_4}{(2\pi)^4}\frac{\partial\mathcal{H}_k^{(4)}(q_1,\cdots,q_4)}{\partial t}\bigg[\tilde{\phi}(q_1)\tilde{\phi}(q_2)\tilde{\phi}(q_3)\tilde{\phi}(q_4) + 3G_k^{(2)}(q_1,q_2)G_k^{(2)}(q_3,q_4) \\ &\qquad + 6G_k^{(2)}(q_1,q_2)\tilde{\phi}(q_3)\tilde{\phi}(q_4) + 4G_k^{(3)}(q_1,q_2,q_3)\tilde{\phi}(q_4) + G_k^{(4)}(q_1,\cdots,q_4)\bigg], \end{aligned} \tag{IV.8}$$

where the full two-point Green's function $G_k^{(2)}(p,q)$ are to be replaced by $G_k^{(2)}(p,q) = G_k^{(2)}(q)\delta(p-q)$ with $G_k^{(2)}(q) = [\Gamma_k^{(2)}(q)]^{-1}$. For the three- and four-point Green's functions, $G_k^{(3)}$ and $G_k^{(4)}$ in the remaining terms of (IV.8), they will be replaced by

$$\begin{aligned} G_k^{(3)}(q_1,q_2,q_3) &= -[\Gamma_k^{(2)}(q_1)]^{-1}[\Gamma_k^{(2)}(q_2)]^{-1}[\Gamma_k^{(2)}(q_3)]^{-1}\Gamma_k^{(3)}(q_1,q_2,q_3). \\ G_k^{(4)}(q_1,\cdots,q_4) &= -[\Gamma_k^{(2)}(q_1)]^{-1}\cdots[\Gamma_k^{(2)}(q_4)]^{-1}\Gamma_k^{(4)}(q_1,q_2,q_3,q_4) \\ &\quad +3\ [\Gamma_k^{(2)}(q_1)]^{-1}\cdots[\Gamma_k^{(2)}(q_4)]^{-1}\int\frac{d^4\ell}{(2\pi)^4}[\Gamma_k^{(2)}(\ell)]^{-1}\Gamma_k^{(3)}(q_1,q_2,\ell)\Gamma_k^{(3)}(\ell,q_3,q_4). \end{aligned} \tag{IV.9}$$

The graphical representation of (IV.8) is demonstrated in Fig. 2. At this stage a couple of remarks are in order. First, let us notice that in the cutoff $\lambda\varphi^4$ theory, the relations between the n-point Green's

---

[2] See also [5, 10].


$\partial_t \Gamma_k[\phi] =$
$\frac{1}{2}$
$\partial_t \mathcal{H}_k^{(2)}$
$+ \frac{1}{2}$
$+$
$+ 3$
$+ 6$
$- 4$
$-$
$+ 3$

FIG. 2: Diagramatic representation of the RG flow equation of the corresponding effective average action for cutoff $\lambda\varphi^4$ theory (IV.8). The thick black line represents full two-point Green's functions $G_k^{(2)} = [\Gamma_k^{(2)}]^{-1}$. The filled red boxes and the small red circles represent the insertion of a factor $\partial_t \mathcal{H}_k^{(2)}$ and $\partial_t \mathcal{H}_k^{(4)}$, respectively. The thin lines connected to ⊠ denote the background field $\phi = \langle\varphi\rangle$. The big gray and black circles are 1PI three- as well as four-point vertex functions, $\Gamma_k^{(3)}$ as well as $\Gamma_k^{(4)}$, respectively.

functions $G_k^{(n)}$ and n-point vertex functions $\Gamma_k^{(n)}$ are not directly affected by the cutoff function $h_k(q)$ [for a proof, see App. C]. This is in contrast to, e.g. (IV.5), where the ordinary relation between two-point Green's function and 1PI two-point vertex function is modified with an additional term including the cutoff function $\mathcal{R}_k(q)$. As it turns out, this is because of the multiplicative nature of $h_k(q)$, in contrast to the additive nature of ΔS_k from (IV.1), consisting of the IR cutoff function $\mathcal{R}_k(q)$. The second point concerns the appearance of new additional contributions in (IV.8) compared to (IV.4). This is because in the cutoff $\lambda\varphi^4$ theory, $h_k(q)$ appears not only in the kinetic part of the classical cut off action, as in the standard derivation of (IV.4), but also in interaction part of the classical action, as all the new contributions are proportional to $\partial_t \mathcal{H}_k^{(4)}$, with $\mathcal{H}_k^{(4)}$ from (IV.7), appearing in the interaction part of S_k from (IV.6). It would be interesting to explore the practical consequences of these new terms in RG flow equation (IV.8). This will be done elsewhere [11]. Let us also note that the procedure leading to (IV.8) can be easily generalized to Abelian and non-Abelian gauge theories. As we have shown in the previous section, a new hidden gauge symmetry associated with the cutoff procedure used in this paper exists, which guarantees the gauge invariance along the flow equation. This is in contrast to the situation of the standard Wetterich's exact RGE, where the manifest gauge invariance is lost, because the regulator is not manifestly gauge invariant [5].

V. CONCLUSIONS

In this work we presented the procedure of introducing a momentum cutoff in a field theory by directly cutting off the momentum on each field via a cutoff function. For QED the resulting exact renormalization group flow equation was shown to respect a generalized "cutoff gauge invariance" which ensures renormalizability, and unitarity without the need for an explicit calculation using modified Ward-Takahashi identities.

We need to emphasize that although the gauge symmetry found is motivated from a noncommutative geometric setup, this symmetry is an inherent symmetry of ordinary QED cut off in momentum, in the spirit of Polchinski's procedure. Our cutoff procedure may also be applied to non-Abelian gauge theories in a similar manner and the resultant exact renormalization group flow be used for calculation of various nonperturbative quantities in QCD [5] and in gravity [12].

There is a subtle point that we would like to comment on in conclusion and that is the singular nature of our cutoff function $h_\Lambda(p)$. As stated earlier this function is quite general, subject only to the restriction that it be equal to unity for momenta smaller than Λ and vanishing rapidly above it. Strictly speaking this function should be chosen to be zero for $p^2 > \Lambda_1^2$, for some $\Lambda_1 > \Lambda$, as in (II.3), and thereby effectively limit the domain of definition of the fields contained in the finite volume of $p^2 < \Lambda_1^2$. This is basically what we have in mind and it makes the inverses of h_Λ well defined, as the domain of definition of fields will then be contained in this volume. In other words, in the expressions in the body of this work, the fields $\varphi(x)$ should be understood as Fourier coefficients of the momentum space fields $\tilde{\varphi}(p)$ which are defined only in the $p^2 < \Lambda_1^2$ region. This also resolves a puzzle that multiplication of the fields by a function h_Λ, if it were smooth, would simply be a field redefinition of the theory and therefore physically trivial. It is also worth to notice that with the sharp cutoff function h_Λ, the theory becomes a modified lattice field theory as the fields are now defined on a discrete set, although the gauge theory would not be the standard lattice gauge theory of Wilson.

For a scale dependent cutoff function, h_k, the RG flow of the effective average action is shown to be different from the standard flow equation from [5, 8–10]. In [11], we will generalize the method leading to the exact RG equation of cutoff $\lambda\varphi^4$ theory to gauge theories and will explore the practical consequences of the new generalized gauge symmetry together with the effect of new terms appearing in the RG flow equation (IV.8) of the effective average action corresponding to these cutoff gauge theories.

VI. ACKNOWLEDGMENTS

The authors thank S. Rouhani, M. Alishahiha and A. E. Mosaffa for useful discussions.

Appendix A: Noncommutative gauge invariance

In this appendix, we review the generalized translationally invariant noncommutative field theory [13, 14], and use the generalized gauge invariance of the relevant noncommutative gauge theory to prove the generalized cutoff gauge invariance in Sec. III, (III.6) and (III.8), of the effective action S_{h_Λ}.

A general star-product of the algebra of functions is the generalization of the usual C^*-algebra of pointwise multiplication algebra of functions. For a translationally invariant product,

$$(f \star g)(x) = \int \frac{d^4p}{(2\pi)^4}\frac{d^4q}{(2\pi)^4}\; e^{ipx}\mathcal{K}(p,q)\tilde{f}(p-q)\tilde{g}(q). \tag{A.1}$$

The point wise multiplication is the special case of $\mathcal{K} = 1$. Associativity of the algebra is the main constraint on the function $\mathcal{K}$. It is

$$\mathcal{K}(p,q)\mathcal{K}(q,r) = \mathcal{K}(p,r)\mathcal{K}(p-r,q-r). \tag{A.2}$$

It was shown in [14] that the following expression is a solution of (A.2)

$$\mathcal{K}(p,q) = h^{-1}(p)h(q)h(p-q)e^{i\Omega(p,q)}, \tag{A.3}$$

with

$$\Omega(p,q) = \theta_{\mu\nu}p^{\mu}q^{\nu} + \eta(q) - \eta(p) + \eta(p-q). \tag{A.4}$$

Here, $h(p)$ is an arbitrary real even and η an arbitrary real odd function. Moreover, $\theta_{\mu\nu}$ is an anti-symmetric constant matrix.

It is readily seen that for $\theta \neq 0$, the algebra is noncommutative, and it is commutative when $\theta = 0$. When $\theta = 0$ and $\eta = 0$, the new star-product in the momentum space, involves multiplications of the functions of the algebra by the fixed function h

$$(f \star g)(x) = \int \frac{d^4p}{(2\pi)^4}\, \frac{1}{h(p)} e^{ipx} \int \frac{d^4q}{(2\pi)^4} [h(p-q)\tilde{f}(p-q)][h(q)\tilde{g}(q)]. \tag{A.5}$$

thus

$$\begin{aligned} h(p)(\widetilde{f \star g})(p) &= \int \frac{d^4q}{(2\pi)^4} [h(p-q)\tilde{f}(p-q)][h(q)\tilde{g}(q)] \\ &= (h\tilde{f}) \circ (h\tilde{g}), \end{aligned} \tag{A.6}$$

where the symbol $\circ$ stands for the convolution product

$$(f \circ g)(x) = \int d^4y\ f(x-y)g(y).$$

In the coordinate space, the product (A.5) could then be defined as

$$(f \star g)(x) = (\widetilde{h^{-1}}) \circ [(\tilde{h} \circ f)(\tilde{h} \circ g)], \tag{A.7}$$

where we have used the relation

$$(\widetilde{fg}) = (\tilde{f} \circ \tilde{g}). \tag{A.8}$$

It must be mentioned that as long as h is a smooth one-to-one map, the new star-algebra is an isomorphism of the algebra with the $C^{\star}$-algebra. However, there is nothing to forbid the function h to be singular. In fact, we have used such a singular h to cut off the momentum of the field theory in the text. In our case the function h effectively cuts down the domain of the function space. The inverse function h^{-1} appearing above should be understood in this context.[3]

The general star-product defined in (A.1) can be used to build a nontrivial "noncommutative field theory", where products of fields in the action are replaced by the new star-product [13, 14]. In particular the action of noncommutative QED would have the form

$$\begin{aligned} S_{\text{QED}} &= \int d^4x \Big[-\bar{\psi}(i\gamma^{\mu}\partial_{\mu} - m) \star \psi + e\bar{\psi} \star \gamma^{\mu}A_{\mu} \star \psi + \frac{1}{4}F_{\mu\nu} \star F^{\mu\nu} + \frac{1}{2\xi}(\partial_{\mu}A^{\mu}) \star (\partial_{\nu}A^{\nu}) \\ &\qquad -\frac{1}{2}\left(i\bar{c} \star \partial^{\mu}D_{\mu}c - i\partial^{\mu}D_{\mu}c \star \bar{c}\right) \Big], \end{aligned} \tag{A.9}$$

---

[3] In this Appendix, we have skipped the subscript Λ in h_{Λ} appearing in Secs. II and III.

where

$$F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu + ie\ (A_\mu \star A_\nu - A_\nu \star A_\mu), \tag{A.10}$$

and $D_\mu c = \partial_\mu c + ie(A_\mu \star\ c - c \star A_\mu)$. This theory is invariant under the star-gauge transformation

$$\psi \to g \star \psi, \qquad \bar{\psi} \to \bar{\psi} \star g^{-1}, \qquad \text{and} \qquad A_\mu \to g \star A \star g^{-1} - \frac{i}{e} g \star \partial_\mu g^{-1}, \tag{A.11}$$

where

$$g = e_\star^{ie\epsilon} = 1 + ie\epsilon + \frac{1}{2!}(ie\epsilon) \star (ie\epsilon) + \cdots . \tag{A.12}$$

See [13, 14] for more details. In this work we have limited ourself to the special case, albeit commutative, of $\theta = 0$, $\eta = 0$ in (A.3).[4] To show the gauge invariance of S_{h_Λ} in Sec. III under the gauge transformations (III.6) and (III.8), it is enough to transform (A.9) into the momentum space using the definition of the star-product (A.5) and

$$\int d^4x (f \star g)(x) = \int \frac{d^4q}{(2\pi)^4} h^2(q) \tilde{f}(-q) \tilde{g}(q), \tag{A.13}$$

where h in (A.5) is identified with the cutoff function h_Λ appearing in (III.2). The resulting expression will be equivalent with S_{h_Λ} arising from (III.1) after the procedure described in (III.2). The gauge transformations (III.6) and (III.8) will then be equivalent with the gauge transformations (A.11) with star-products defined in (A.7).

Appendix B: The Proof of Ward identitiies

To prove the Ward identities of cutoff QED, we start with the generating functional for the full Green's functions

$$Z[J_\mu, \chi, \bar{\chi}] = \int \mathcal{D}\psi\ \mathcal{D}\bar{\psi}\ \mathcal{D}A_\mu\ e^{iS_{\text{tot}}}, \tag{B.1}$$

with $S_{\text{tot}} = S_{\text{QED}} + S_{\text{source}}$. Here, S_{QED} is given by (A.9), where the ghost terms can be ignored, and S_{source} by

$$S_{\text{source}} = \int d^4x \left(\bar{\chi} \star \psi + \bar{\psi} \star \chi + J_\mu \star A^\mu \right). \tag{B.2}$$

Varying $Z[J_\mu, \chi, \bar{\chi}]$ in (B.1) with respect to the star-gauge transformation (A.11) and replacing the star-products with the expression on the r.h.s. of (A.13) to introduce the cutoff function h_Λ, we arrive at

$$\begin{aligned} \Big\{ &- \frac{1}{\xi} p^2 p^\mu h_\Lambda^2(p) \tilde{A}_\mu(p) + p^\mu h_\Lambda^2(p) \tilde{J}_\mu(p) \\ &-e h_\Lambda(p) \int \frac{d^4q}{(2\pi)^4} \Big[h_\Lambda(p-q) \tilde{\bar{\psi}}(q-p) h_\Lambda(q) \tilde{\chi}(q) - h_\Lambda(p-q) \tilde{\bar{\chi}}(q-p) h_\Lambda(q) \tilde{\psi}(q) \Big] \Big\} Z = 0. \end{aligned} \tag{B.3}$$

---

[4] With this special choice of θ and η, the field strength tensor (A.10) turns out to be the same as in commutative QED. The ghosts will also decouple from the theory.

Using the relations[5]

$$\tilde{\psi}(p) = h_{\Lambda}^{-2}(p)\frac{\delta Z}{i\delta\tilde{\bar{\chi}}(p)}, \qquad \tilde{\bar{\psi}}(p) = -h_{\Lambda}^{-2}(p)\frac{\delta Z}{i\delta\tilde{\chi}(p)}, \qquad \tilde{A}_{\mu}(p) = h_{\Lambda}^{-2}(p)\frac{\delta Z}{i\delta\tilde{J}_{\mu}(-p)}, \tag{B.4}$$

and replacing $Z[J_\mu, \chi, \bar{\chi}]$ by $Z = e^{iW}$, with $W[J_\mu, \chi, \bar{\chi}]$ the generating function of connected Green's functions, we arrive at

$$\begin{aligned}
&-\frac{1}{\xi}p^2 p_\mu \frac{\delta W}{\delta \tilde{J}_\mu(-p)} + h_\Lambda^2(p)p_\mu \tilde{J}^\mu(p)\\
&\quad + e h_\Lambda(p)\int\frac{d^4q}{(2\pi)^4}\left[h_\Lambda(q)h_\Lambda^{-1}(q-p)\frac{\delta W}{\delta\tilde{\chi}(q-p)}\tilde{\chi}(q) + h_\Lambda^{-1}(q)h_\Lambda(q-p)\tilde{\bar{\chi}}(q-p)\frac{\delta W}{\delta\tilde{\bar{\chi}}(q)}\right] = 0.
\end{aligned} \tag{B.5}$$

To derive (B.4), $h_\Lambda(p) = h_\Lambda(-p)$ is used. To arrive at the Ward identity in terms of $\Gamma[A_\mu, \psi, \bar{\psi}]$, the generating functional for 1PI Green's function, we use the Legendre transformation (III.11) leading to

$$\begin{aligned}
\frac{\delta W}{\delta \tilde{J}^\mu(-p)} &= h_\Lambda^2(p)\tilde{A}_\mu(p), & \frac{\delta W}{\delta\tilde{\chi}(p)} &= -h_\Lambda^2(p)\tilde{\bar{\psi}}(p), & \frac{\delta W}{\delta\tilde{\bar{\chi}}(p)} &= h_\Lambda^2(p)\tilde{\psi}(p),\\
\tilde{J}_\mu(p) &= -h_\Lambda^{-2}(p)\frac{\delta\Gamma}{\delta\tilde{A}^\mu(-p)}, & \tilde{\chi}(p) &= -h_\Lambda^{-2}(p)\frac{\delta\Gamma}{\delta\tilde{\bar{\psi}}(p)}, & \tilde{\bar{\chi}}(p) &= h_\Lambda^{-2}(p)\frac{\delta\Gamma}{\delta\tilde{\psi}(p)}.
\end{aligned} \tag{B.6}$$

Plugging these relations in (B.5), we arrive at

$$\begin{aligned}
&-\frac{1}{\xi}p^2 p_\mu h_\Lambda^2(p)A^\mu(p) - p_\mu\frac{\delta\Gamma}{\delta\tilde{A}_\mu(-p)}\\
&\quad + e h_\Lambda(p)\int\frac{d^4q}{(2\pi)^4}\left[h_\Lambda^{-1}(q)h_\Lambda(q-p)\tilde{\bar{\psi}}(q-p)\frac{\delta\Gamma}{\delta\tilde{\bar{\psi}}(q)} + h_\Lambda(q)h_\Lambda^{-1}(q-p)\frac{\delta\Gamma}{\delta\tilde{\psi}(q-p)}\tilde{\psi}(q)\right] = 0.
\end{aligned} \tag{B.7}$$

As a first example on the application of (B.5), let us differentiate it with respect to $\tilde{J}_\nu(p')$ and set eventually $\tilde{J}_\mu = \tilde{\bar{\chi}} = \tilde{\chi} = 0$. We arrive at the Ward identity for the full photon propagator of the cutoff QED in momentum space, $\tilde{D}_\Lambda^{\mu\nu}$,

$$\frac{i}{\xi}p^2 p_\mu h_\Lambda^2(p)\tilde{D}_\Lambda^{\mu\nu}(p) = p^\nu. \tag{B.8}$$

Here, we have used (B.6) to get first

$$\left.\frac{\delta W}{\delta\tilde{J}_\mu(-p)\delta\tilde{J}_\nu(p')}\right|_{\tilde{J}_\mu=\tilde{\bar{\chi}}=\tilde{\chi}=0} = h_\Lambda^4(p)\langle\tilde{A}^\mu(p)\tilde{A}^\nu(-p')\rangle, \tag{B.9}$$

and defined the full cutoff dependent photon propagator $\tilde{D}_\Lambda^{\mu\nu}(p)$ by

$$i\tilde{D}_\Lambda^{\mu\nu}(p)\delta(p-p') \equiv \langle\tilde{A}^\mu(p)\tilde{A}^\nu(-p')\rangle. \tag{B.10}$$

Equation (B.8) is in particular satisfied by tree level photon propagator of cutoff QED [14]

$$\left.\tilde{D}_\Lambda^{\mu\nu}(p)\right|_{\text{tree-level}} = -\frac{ih_\Lambda^{-2}(p)}{p^2}\left(g^{\mu\nu} - (1-\xi)\frac{p^\mu p^\nu}{p^2}\right). \tag{B.11}$$

---

[5] We follow the notations in [15].

As a second example, let us differentiate (B.7) with respect to $\tilde{\psi}(-\ell)$ and $\tilde{\bar{\psi}}(k)$ and set eventually $\tilde{A}_\mu = \tilde{\psi} = \tilde{\bar{\psi}} = 0$. Using (B.6), we arrive at

$$\frac{1}{h_\Lambda(p)h_\Lambda(\ell+p)h_\Lambda(\ell)} p_\mu \tilde{\Gamma}^\mu_\Lambda(-\ell-p,-\ell;-p) \;=\; e\left[h_\Lambda^{-2}(\ell+p)\tilde{S}_\Lambda^{-1}(\ell+p) - h_\Lambda^{-2}(\ell)\tilde{S}_\Lambda^{-1}(\ell)\right], \tag{B.12}$$

where the 1PI three point vertex function

$$\tilde{\Gamma}^\mu_\Lambda(k,-\ell;-p)\delta(k+p+\ell) \equiv \frac{\delta^3\Gamma}{\delta\tilde{\bar{\psi}}(k)\delta\tilde{\psi}(-\ell)\delta\tilde{A}_\mu(-p)}\Bigg|_{\tilde{A}_\mu=\tilde{\psi}=\tilde{\bar{\psi}}=0}, \tag{B.13}$$

as well the fermionic 1PI two-point function at finite cutoff Λ

$$\tilde{S}_\Lambda^{-1}(\ell)\delta(k+\ell) \equiv \frac{\delta^2\Gamma}{\delta\tilde{\bar{\psi}}(k)\delta\tilde{\psi}(-\ell)}\Bigg|_{\tilde{\psi}=\tilde{\bar{\psi}}=0}, \tag{B.14}$$

are introduced. Taking the limit $p \to 0$ in (B.12) and using $h_\Lambda(0) = 1$, we arrive at the standard relation

$$\tilde{\Gamma}^\mu_\Lambda(-\ell,-\ell;0) = e\frac{\partial\tilde{S}_\Lambda^{-1}(\ell)}{\partial\ell_\mu}. \tag{B.15}$$

Assuming that $h_\Lambda(p)$ is a nearly constant function for $|p| < \Lambda$, and using[6]

$$\begin{aligned}\tilde{S}_\Lambda^{-1}(\ell) \;&=\; h_\Lambda^2(\ell)\tilde{S}_\infty^{-1}(\ell),\\ \tilde{\Gamma}^\mu_\Lambda(k,\ell;p) \;&=\; h_\Lambda(k)h_\Lambda(\ell)h_\Lambda(p)\tilde{\Gamma}_\infty(k,\ell;p),\end{aligned} \tag{B.16}$$

we get

$$\tilde{\Gamma}^\mu_\infty(-\ell,-\ell;0) = e\frac{\partial\tilde{S}_\infty^{-1}(\ell)}{\partial\ell_\mu}, \tag{B.17}$$

where $\tilde{S}_\infty$ and $\tilde{\Gamma}^\mu_\infty$ are the 1PI two- and three-point ertex functions of QED in $\Lambda \to \infty$ limit. Assuming at this stage that (B.17) is also valid for renormalized Green's functions $[\tilde{\Gamma}^\mu_\infty]_r = Z_2^{-1}Z_3^{-1/2}\tilde{\Gamma}^\mu_\infty$ and $[\tilde{S}_\infty^{-1}]_r = Z_2^{-1}\tilde{S}_\infty^{-1}$, as well as for renormalized coupling $e_r \equiv Z_1 Z_2^{-1} Z_3^{-1/2} e$, with Z_1, Z_2 and Z_3 the renormalization constants corresponding to the vertex function, fermion and photon propagators, respectively, we arrive at $Z_1 = Z_2$.

Appendix C: Exact RG flow equation of cutoff $\lambda\varphi^4$ theory

Let us start by considering the bare action of $\lambda\varphi^4$ theory in Euclidean space (II.1). As we have explained in Sec. IV, each field $\tilde{\varphi}(q)$ shall be replaced by $h_k(q)\varphi(q)$, where k is the renormalization scale. The modified classical action is then given by (IV.6) with the cutoff functions given in (IV.7). The corresponding generating functional of this cutoff theory then reads

$$Z_k[J] = \int \mathcal{D}\varphi\ \exp\left(-S_k[\varphi] + \int \frac{d^4q}{(2\pi)^4} h_k^2(q)\tilde{J}(-q)\tilde{\varphi}(q)\right), \tag{C.1}$$

---

[6] Relations (B.16) are shown to be valid at one-loop level (see [14] for more details).

where $h_k(-q) = h_k(q)$ is assumed. The Legendre transformation between $W_k[J] \equiv \ln Z_k[J]$, the generating functional of the connected Green's function, and the 1PI effective average action, $\Gamma_k[\phi]$ is given by

$$\Gamma_k[\phi] = -W_k[J] + \int \frac{d^4q}{(2\pi)^4} h_k^2(q)\tilde{J}(-q)\tilde{\phi}(q), \tag{C.2}$$

where $\phi \equiv \langle\varphi\rangle$, and

$$\frac{\delta W_k[J]}{\delta \tilde{J}(-q)} = h_k^2(q)\tilde{\phi}(q). \tag{C.3}$$

It is the purpose of this appendix to derive the scale dependence of $\Gamma_k[\phi]$. To do this, we will follow the method described in [5, 8, 10]. First, we differentiate $\Gamma_k[\phi]$ from (C.2) with respect to k and arrive at

$$\frac{\partial \Gamma_k[\phi]}{\partial k} = -\frac{\partial W_k[J]}{\partial k} + 2\int \frac{d^4q}{(2\pi)^4} h_k(q)\frac{\partial h_k(q)}{\partial k}\tilde{J}(-q)\tilde{\phi}(q) = \left\langle \frac{\partial S_k[\varphi]}{\partial k}\right\rangle, \tag{C.4}$$

where (C.3) and the standard notation

$$\langle \mathcal{O}[\varphi]\rangle = Z_k^{-1}[J]\int \mathcal{D}\varphi\ \mathcal{O}[\varphi]\ \exp\left(-S_k[\varphi] + \int \frac{d^4q}{(2\pi)^4} h_k^2(q)\tilde{J}(-q)\tilde{\varphi}(q)\right), \tag{C.5}$$

are used. Plugging now (IV.6) in (C.4), we get

$$\begin{aligned}\frac{\partial \Gamma_k[\phi]}{\partial k} = \left\langle \frac{\partial S_k[\varphi]}{\partial k}\right\rangle &= \frac{1}{2}\int \frac{d^4q_1}{(2\pi)^4}\frac{d^4q_2}{(2\pi)^4}\frac{\partial \mathcal{H}_k^{(2)}(q_1,q_2)}{\partial k}\langle\tilde{\varphi}(q_1)\tilde{\varphi}(q_2)\rangle \\ &+ \int \frac{d^4q_1}{(2\pi)^4}\cdots\frac{d^4q_4}{(2\pi)^4}\frac{\partial \mathcal{H}_k^{(4)}(q_1,\cdots,q_4)}{\partial k}\langle\tilde{\varphi}(q_1)\tilde{\varphi}(q_2)\tilde{\varphi}(q_3)\tilde{\varphi}(q_4)\rangle.\end{aligned} \tag{C.6}$$

At this stage we shall replace the two- and four-point Green's functions appearing on the r.h.s. of (C.6) by a combination of connected and disconnected Green's functions. To do this, let us vary $W_k[J] = \ln Z_k[J]$ two times with respect to $\tilde{J}$ to get

$$\begin{aligned}\frac{\delta^2 W_k[J]}{\delta\tilde{J}(-q_1)\delta\tilde{J}(-q_2)} &= -\frac{1}{Z_k^2[J]}\frac{\delta Z_k[J]}{\delta\tilde{J}(-q_1)}\frac{\delta Z_k[J]}{\delta\tilde{J}(-q_2)} + \frac{1}{Z_k[J]}\frac{\delta^2 Z_k[J]}{\delta\tilde{J}(-q_1)\delta\tilde{J}(-q_2)} \\ &= h_k^2(q_1)h_k^2(q_2)\left(-\tilde{\phi}(q_1)\tilde{\phi}(q_2) + \langle\tilde{\varphi}(q_1)\tilde{\varphi}(q_2)\rangle\right).\end{aligned} \tag{C.7}$$

Here, the definition of $Z_k[J]$ from (C.1) is used. Defining then the connected n-point Green's functions as

$$\frac{\delta^n W_k[J]}{\delta\tilde{J}(-q_1)\cdots\tilde{J}(-q_n)} \equiv h_k^2(q_1)\cdots h_k^2(q_n)G_k^{(n)}(q_1,\cdots,q_n), \tag{C.8}$$

and plugging the corresponding relation for $n=2$ on the l.h.s. of (C.7), we get the standard relation

$$\langle\tilde{\varphi}(q_1)\tilde{\varphi}(q_2)\rangle = G_k^{(2)}(q_1,q_2) + \tilde{\phi}(q_1)\tilde{\phi}(q_2). \tag{C.9}$$

Similar relation exists also between the connected four-point Green's function $G_k^{(4)}(q_1,\cdots,q_4)$ and $\langle\tilde{\varphi}(q_1)\tilde{\varphi}(q_2)\tilde{\varphi}(q_3)\tilde{\varphi}(q_4)\rangle$ appearing on the r.h.s. of (C.6). It is given by

$$\begin{aligned}\langle\tilde{\varphi}(q_1)\tilde{\varphi}(q_2)\tilde{\varphi}(q_3)\tilde{\varphi}(q_4)\rangle &= G_k^{(4)}(q_1,\cdots,q_4)\\ &+G_k^{(3)}(q_1,q_2,q_3)\tilde{\phi}(q_4)+G_k^{(3)}(q_1,q_2,q_4)\tilde{\phi}(q_3)+G_k^{(3)}(q_1,q_4,q_3)\tilde{\phi}(q_2)+G_k^{(3)}(q_4,q_2,q_3)\tilde{\phi}(q_1)\\ &+G_k^2(q_1,q_2)\tilde{\phi}(q_3)\tilde{\phi}(q_4)+G_k^2(q_1,q_3)\tilde{\phi}(q_2)\tilde{\phi}(q_4)+G_k^2(q_1,q_4)\tilde{\phi}(q_2)\tilde{\phi}(q_3)+G_k^2(q_2,q_3)\tilde{\phi}(q_1)\tilde{\phi}(q_4)\\ &+G_k^2(q_2,q_4)\tilde{\phi}(q_1)\tilde{\phi}(q_3)+G_k^2(q_3,q_4)\tilde{\phi}(q_1)\tilde{\phi}(q_2)\\ &+G_k^{(2)}(q_1,q_2)G_k^{(2)}(q_3,q_4)+G_k^{(2)}(q_1,q_3)G_k^{(2)}(q_2,q_4)+G_k^{(2)}(q_1,q_4)G_k^{(2)}(q_2,q_3)\\ &+\tilde{\phi}(q_1)\tilde{\phi}(q_2)\tilde{\phi}(q_3)\tilde{\phi}(q_4).\end{aligned}\tag{C.10}$$

Plugging (C.9) and (C.10) in (C.6) and using the symmetry of $\mathcal{H}_k^{(2)}(q_1,q_2)$ and $\mathcal{H}_k^{(4)}(q_1,\cdots,q_4)$ under permutation of $q_i, i=1,\cdots,4$, we arrive at the flow equation of $\Gamma_k[\phi]$ in terms of the connected $n=1,\cdots,4$-point Green's functions

$$\begin{aligned}\frac{\partial\Gamma_k[\phi]}{\partial t} &= \frac{1}{2}\int\frac{d^4q_1}{(2\pi)^4}\frac{d^4q_2}{(2\pi)^4}\frac{\partial\mathcal{H}_k^{(2)}(q_1,q_2)}{\partial t}[G_k^{(2)}(q_1,q_2)+\tilde{\phi}(q_1)\tilde{\phi}(q_2)]\\ &+\int\frac{d^4q_1}{(2\pi)^4}\cdots\frac{d^4q_4}{(2\pi)^4}\frac{\partial\mathcal{H}_k^{(4)}(q_1,\cdots,q_4)}{\partial t}\bigg[\tilde{\phi}(q_1)\tilde{\phi}(q_2)\tilde{\phi}(q_3)\tilde{\phi}(q_4)+3G_k^{(2)}(q_1,q_2)G_k^{(2)}(q_3,q_4)\\ &\qquad+6G_k^{(2)}(q_1,q_2)\tilde{\phi}(q_3)\tilde{\phi}(q_4)+4G_k^{(3)}(q_1,q_2,q_3)\tilde{\phi}(q_4)+G_k^{(4)}(q_1,\cdots,q_4)\bigg].\end{aligned}\tag{C.11}$$

This flow equation is to be compared with the standard flow equation (IV.4), where only a term similar to the first term on the r.h.s. of (C.11) appears. The appearance of additional terms in (C.11), including the contributions of $G_k^{(n)}(q_1,\cdots,q_n), n=1,\cdots,4$, is, in particular, a consequence of the replacement of $\tilde{\varphi}(q)$ by $h_k(q)\tilde{\varphi}(q)$ in the interaction term of the original classical action, in contrast to the standard procedure [5, 8–10].

In a last step, we shall use the relations between the connected n-point Green's function, $G_k^{(n)}(q_1,\cdots,q_n)$, defined in (C.8) and the 1PI n-point vertex functions, $\Gamma_k^{(n)}(q_1,\cdots,q_n)$, defined by

$$\Gamma_k^{(n)}(q_1,\cdots,q_n)\equiv\frac{\delta^2\Gamma_k[\phi]}{\delta\tilde{\phi}(q_1)\cdots\delta\tilde{\phi}(q_n)},\tag{C.12}$$

to replace $G_k^{(n)}(q_1,\cdots,q_n), n=1,\cdots,4$ in (C.11) by the corresponding expressions in terms of $\Gamma_k^{(n)}(q_1,\cdots,q_n)$. To do this, let us first consider the relation

$$\int\frac{d^4\ell_1}{(2\pi)^4}\frac{\delta\tilde{J}(-q_1)}{\delta\tilde{\phi}(\ell_1)}\frac{\delta\tilde{\phi}(\ell_1)}{\delta\tilde{J}(-q_2)}=\delta(q_1-q_2).\tag{C.13}$$

It can easily be shown that

$$\frac{\delta\tilde{J}(-q_1)}{\delta\tilde{\phi}(\ell_1)}=h_k^{-2}(q_1)\Gamma_k^{(2)}(q_1,\ell_1),\qquad\text{and}\qquad\frac{\delta\tilde{\phi}(\ell_1)}{\delta\tilde{J}(-q_2)}=h_k^2(q_2)G_k^{(2)}(\ell_1,q_2).\tag{C.14}$$

Plugging (C.14) in (C.13), and using $G_k^{(2)}(\ell_1,q_2)=G_k^{(2)}(q_2)\delta(\ell_1-q_2)$ as well as $\Gamma_k^{(2)}(q_1,\ell_1)=\Gamma_k^{(2)}(q_1)\delta(q_1-\ell_1)$, we arrive, after integrating over ℓ_1, at

$$G_k^{(2)}(q)\Gamma_k^{(2)}(q)=1.\tag{C.15}$$

This is the standard relation between $G_k^{(2)}(q)$ and $\Gamma_k^{(2)}(q)$. In contrast to (IV.5), the cutoff function $h_k(q)$ does not appear in (C.15). This is because of the multiplicative nature of the cutoff function $h_k(q)$. Similar relation can also be derived between $G_k^{(3)}$ and $\Gamma_k^{(3)}$. It is simplify given by

$$G_k^{(3)}(q_1,q_2,q_3) = -[\Gamma_k^{(2)}(q_1)]^{-1}[\Gamma_k^{(2)}(q_2)]^{-1}[\Gamma_k^{(2)}(q_3)]^{-1}\Gamma_k^{(3)}(q_1,q_2,q_3), \tag{C.16}$$

which is derived by differentiating (C.13) with respect to $\tilde{J}(-q_3)$, and plugging (C.14) as well as

$$\begin{aligned}
\frac{\delta^2\tilde{J}(-q_1)}{\delta\tilde{\phi}(\ell_1)\delta\tilde{J}(-q_3)} &= h_k^{-2}(q_1)h_k^2(q_3)\int\frac{d^4\ell_2}{(2\pi)^4}G_k^{(2)}(\ell_2,q_3)\Gamma_k^{(3)}(q_1,\ell_1,\ell_2),\\
\frac{\delta^2\tilde{\phi}(\ell_1)}{\delta\tilde{J}(-q_2)\delta\tilde{J}(-q_3)} &= h_k^2(q_2)h_k^2(q_3)G_k^{(3)}(\ell_1,q_2,q_3).
\end{aligned} \tag{C.17}$$

in the resulting expression. Similarly, to determine the relation between $G_k^{(4)}$ and $\Gamma_k^{(4)}$, we differentiate (C.13) with respect to $\tilde{J}(-q_3)$ and $\tilde{J}(-q_4)$. Plugging (C.14), (C.17) and

$$\begin{aligned}
\frac{\delta^3\tilde{J}(-q_1)}{\delta\tilde{\phi}(\ell_1)\delta\tilde{J}(-q_3)\delta\tilde{J}(-q_4)} &= h_k^{-2}(q_1)h_k^2(q_3)h_k^2(q_4)\bigg\{\int\frac{d^4\ell_2}{(2\pi)^4}G_k^{(3)}(\ell_2,q_3,q_4)\Gamma_k^{(3)}(q_1,\ell_1,\ell_2)\\
&\quad+\int\frac{d^4\ell_2}{(2\pi)^4}\frac{d^4\ell_3}{(2\pi)^4}G_k^{(2)}(\ell_2,q_3)G_k^{(2)}(\ell_3,q_4)\Gamma_k^{(4)}(q_1,\ell_1,\ell_2,\ell_3)\bigg\},\\
\frac{\delta^3\tilde{\phi}(\ell_1)}{\delta\tilde{J}(-q_2)\delta\tilde{J}(-q_3)\delta\tilde{J}(-q_4)} &= h_k^2(q_2)h_k^2(q_3)h_k^2(q_4)G_k^{(4)}(\ell_1,q_2,q_3,q_4),
\end{aligned} \tag{C.18}$$

in the resulting expression, we arrive after some algebra at

$$\begin{aligned}
G_k^{(4)}(q_1,\cdots,q_4) &= -[\Gamma_k^{(2)}(q_1)]^{-1}\cdots[\Gamma_k^{(2)}(q_4)]^{-1}\bigg\{\Gamma_k^{(4)}(q_1,q_2,q_3,q_4)\\
&-\int\frac{d^4\ell}{(2\pi)^4}[\Gamma_k^{(2)}(\ell)]^{-1}\bigg[\Gamma_k^{(3)}(q_1,q_2,\ell)\Gamma_k^{(3)}(\ell,q_3,q_4)+\Gamma_k^{(3)}(q_1,q_3,\ell)\Gamma_k^{(3)}(\ell,q_2,q_4)\\
&+\Gamma_k^{(3)}(q_1,q_4,\ell)\Gamma_k^{(3)}(\ell,q_2,q_3)\bigg]\bigg\},
\end{aligned} \tag{C.19}$$

where (C.15) and (C.16) are also used. Plugging (C.15), (C.16) and (C.19) in the flow equation (C.11) and using the symmetry of the cutoff function $\mathcal{H}_k^{(4)}(q_1,\cdots,q_4)$ under permutation of $q_i, i=1,\cdots,4$, we arrive at (IV.8).

---

[1] N. N. Bogoliubov, O. S. Parasiuk, *On the multiplication of the causal function in the quantum theory of fields*, Acta Math. **97**, 227 (1957).

K. Hepp, *Proof of the Bogoliubov-Parasiuk theorem on renormalization*, Commun. Math. Phys. **2**, 301 (1966).

Y. Hahn and W. Zimmermann, *An elementary proof of Dyson's power counting theorem*, Comm. Math. Phys. **10**, 330 (1968).

W. Zimmermann, *The power counting theorem for Minkowski metric*, Commun. Math. Phys. **11**, 1 (1968).

[2] J. Polchinski, *Renormalization and effective Lagrangians*, Nucl. Phys. **B231**, 269 (1984).

[3] G. Keller and C. Kopper, *Perturbative renormalization of QED via flow equation*, Phys. Lett. **B273**, 323 (1991); *ibid.*, *Renormalizability proof for QED based on flow equations*, Commun. Math. Phys. **176**, 193 (1996).

H. Sonoda, *On the construction of QED using ERG*, J. Phys. **A40**, 9675 (2007), hep-th/0703167.

[4] B. J. Warr, *Renormalization of gauge theories using effective Lagrangians. 1.*, Annals Phys. **183**, 1 (1988); *ibid.*, *Renormalization of gauge theories using effective Lagrangians. 2.*, Annals Phys. **183**, 59 (1988).

M. Bonini, M. D'Attanasio, G. Marchesini, *Ward identities and Wilson renormalization group for QED*, Nucl. Phys. **B418**, 81 (1994), hep-th/9307174; *ibid.*, *Renormalization group flow for SU(2) Yang-Mills theory and gauge invariance*, Nucl. Phys. **B421**, 429 (1994), hep-th/9312114.

M. Reuter, C. Wetterich, *Effective average action for gauge theories and exact evolution equations*, Nucl. Phys. **B417**, 181 (1994); *ibid.*, *Exact evolution equation for scalar electrodynamics*, Nucl. Phys. **B427**, 291 (1994).

U. Ellwanger, *Flow equations and BRS invariance for Yang-Mills theories*, Phys. Lett. **B335**, 364 (1994), hep-th/9402077.

T. R. Morris, *A Manifestly gauge invariant exact renormalization group*, hep-th/9810104.

M. Simionato, *Gauge consistent Wilson renormalization group: Abelian case*, Int. J. Mod. Phys. **A15**, 2121 (2000), hep-th/9809004; *ibid.* M. Simionato, *Gauge consistent Wilson renormalization group. 2. Non-Abelian case*, Int. J. Mod. Phys. **A15**, 2153 (2000), hep-th/9810117.

V. Branchina, K. A. Meissner, G. Veneziano, *The price of an exact, gauge invariant RG flow equation*, Phys. Lett. **B574**, 319 (2003), hep-th/0309234.

O. J. Rosten, *A Manifestly gauge invariant and universal calculus for SU(N) Yang-Mills*, Int. J. Mod. Phys. **A21**, 4627 (2006), hep-th/0602229.

[5] J. M. Pawlowski, *Aspects of the functional renormalisation group*, Annals Phys. **322**, 2831 (2007), hep-th/0512261.

H. Gies, *Introduction to the functional RG and applications to gauge theories*, hep-ph/0611146.

[6] K. G. Wilson, J. B. Kogut, *The renormalization group and the epsilon expansion*, Phys. Rept. **12**, 75-200 (1974).

[7] C. Wetterich, *Average action and the renormalization group equations*, Nucl. Phys. **B352**, 529-584 (1991).

[8] C. Wetterich, *The average action for scalar fields near phase transitions*, Z. Phys. **C57**, 451 (1993); *ibid. Improvement of the average action,* Z. Phys. **C60**, 461 (1993).

[9] T. R. Morris, *The exact renormalization group and approximate solutions*, Int. J. Mod. Phys. **A9**, 2411 (1994), hep-ph/9308265.

[10] J. Berges, N. Tetradis, C. Wetterich, *Nonperturbative renormalization flow in quantum field theory and statistical physics*, Phys. Rept. **363**, 223 (2002), hep-ph/0005122.

[11] F. Ardalan and N. Sadooghi et al., work in progress.

[12] M. Reuter, *Nonperturbative evolution equation for quantum gravity*, Phys. Rev. **D57**, 971 (1998), hep-th/9605030.

D. F. Litim, *Fixed points of quantum gravity*, Phys. Rev. Lett. **92**, 201301 (2004), hep-th/0312114.

[13] S. Galluccio, F. Lizzi and P. Vitale, *Translation invariance, commutation relations and ultraviolet/infrared mixing,*" JHEP **0909**, 054 (2009), arXiv: 0907.3640 [hep-th].

A. Tanasa and P. Vitale, *Curing the UV/IR mixing for field theories with translation-invariant $\star$ products*, Phys. Rev. D **81**, 065008 (2010), arXiv: 0912.0200 [hep-th].

A. Tanasa, *Translation-invariant noncommutative renormalization*, SIGMA **6**, 047 (2010), arXiv: 1003.4877 [hep-th].

[14] F. Ardalan, N. Sadooghi, *Translational-invariant noncommutative gauge theory*, Phys. Rev. **D83**, 025014 (2011), arXiv: 1008.5064 [hep-th].

[15] S. Pokorski, *Gauge Field Theory*, Second Edition, Cambridge University Press (2001).